

SECTION II

2011

FIRST AID

RULES

2011 FIRST AID CONTEST RULES

INDEX

<u>Title</u>	<u>Section II</u>	<u>Page</u>
Rules Governing the 2011 First Aid Contest.....		1
Scorecard A Discounts		7
Interpretations of Scorecard B (Artificial Ventilation/Cardiopulmonary Resuscitation).		11
<u>Skill Sheets</u>		
Initial Assessment		14
Patient Assessment		15
One-Person CPR (Manikin Only)		17
Two-Rescuer CPR (Manikin Only).....		19
Mouth-to-Mask Resuscitation		23
Airway Obstruction		24
Sucking Chest Wound.....		28
Life Threatening Bleeding.....		29
Tourniquet.....		30
Dressings and Bandaging – Open Wounds		31
Two-Person Log Roll		33
Splinting		34
One Rescuer Blanket Drag.....		42
Ground Lifts.....		43
Shirt Drag		44
Establishing Airway – Suspected Cervical Spine (Neck) Injury		45
Shock.....		46
Immobilization		46
Burns		50
Frostbite.....		52
Late or Deep Cold Injury		53
Hyperthermia		54

RULES GOVERNING THE 2011 NATIONAL FIRST AID CONTEST

First Aid rules were designed as a training tool for first aid teams. They were developed for contest purposes only. Discretion should be used in actual mine emergency situations.

1. Members of First Aid Teams must be bona fide employees of the mining industry. Teams shall furnish their own recording manikin, and all other materials listed in G Miscellaneous.
2. A team shall consist of two members and a patient. A team shall not use the same patient for multiple teams. Bystander(s) may only assist in supporting, lifting, or moving the patient. If the problem requires a bystander(s), they will be provided. Bystander(s) will be positioned at the field and will be identified as a bystander with labels or name tags and BSI precautions will be in place. A team shall work two problems and the scores shall be combined to determine the team's final standings.
3. Each team entering the contest will draw a number to determine the order of the performance at the time of registration. Teams sharing equipment must notify the registrar upon registration, so that these teams will not be working at the same time and enough time is allowed for inspection of equipment prior to second team using the equipment.
4. Each participating team must be under guard before the start of the contest. Any team or team member receiving information concerning a contest problem prior to arriving at the working area will, be disqualified by the Chief Judge and Director.
5. No practicing will be allowed on the field before the beginning of the Contest. No reference books or training material will be permitted in the working area during the working or reading of the problems.
6. Only designated officials will be allowed to communicate with teams while teams are working.
7. All Procedures shall be performed in the order listed: Skill sheets supersede First Aid Rules which supersede Brady First Responder eighth Edition by Bergeron and Le Baudour.
8. Contest officials will designate spaces (15 feet by 15 feet - minimum) for teams to work. All equipment (except manikin) will be kept behind a

baseline designated by a Contest Official. All problems will be worked in the designated area which shall contain only the judges and the contesting teams. The manikin will be located within the designated space.

9. All team members and patient shall be dressed alike. Shoes need not be identical. The patient may wear shorts even if the team members are wearing pants. The pants and shorts shall be the same color.
10. The Timekeepers shall explain to the team the timing devices used. Judges will require a signature and team number on the sample CPR tape at this time. (Check shallow breaths, deep and shallow compressions.)
11. Problems will be kept in unsealed envelopes, retained by the judges, and given to the team after the timing device has been started. Judges shall place the patient in the required position as stated in the problem to be worked.
 - a. The team's material and equipment (jump kits, splints, etc.) may not be assembled or donned (excluding BSI) until after the timing device is started. The manikin may be placed in the designated area prior to starting the timing device.
 - b. The working time for a problem will start when the team starts the timing device.
 - c. If props are to be utilized during the working of the problem, such props must be readily available to the working teams and in working condition. These props must be identified by the judges to the team members prior to starting the timing device and must be located within the designated working area. Props will not be utilized in lieu of first aid equipment for treatment of patient(s). Props will be limited to items related to communication and mechanism of injury for effects unless skill sheets are provided. Props shall be within the application of the skill sheets used for treatment of the injury/conditions.

A barrier device must be used when contacting manikin. The face masks/shields may be removed when the team is required to give artificial ventilation, CPR, inflating splints, etc.

12. Injuries/conditions requiring treatment will be identified by cards, envelopes or labels attached to the patient at or as near the location of the injury as possible on the outside of the clothing, be identified by simulated wounds, or be in the reading of the problem. Signs, symptoms or mechanisms of injury

may be used. If signs and symptoms are used, all signs and symptoms shall be identified by cards, envelopes or labels placed on patient. All signs and symptoms will be given to the teams in writing. Wounds that are listed in the reading of the problem shall also be placed on patient. (Exception: If the wound is on the eyelid or an impaled object in the eye, the label will NOT be placed on the eye, but in an obvious area near the eye.)

During the initial or patient assessment, teams may find an envelope attached to the patient or be provided an envelope by the judges which contains patient information that needs immediate attention. If repositioning of patient is required for treatment, patient must be placed in the proper position prior to treatment. Upon completion of treatment of these conditions, the initial or patient assessment will be resumed at the point where team left off. The patient will already be marked upon arrival of the team.

Since spinal stabilization must be taken as outlined in Initial Assessment, spinal injuries will be stated in the reading of the problem and not marked on the patient. Sucking chest wounds and life-threatening bleeding shall be stated in the reading of the problem and shall be marked on the patient.

13. Lettering on the cards and/or labels will be at least 1/4-inch in height.

Example:

2-INCH WOUND ON FOREHEAD

14. The problem will end and teams will stop the timing device when all conditions have been located, treated, and work area has been cleaned. The timekeeper/judge must time the problem in minutes and seconds and consult with the team upon completion of the problem to verify the time.
15. After stopping the timing device, the team will inform the judges that the patient is ready for inspection. Team members will remain with the patient until released by the judges. Any physical treatment(s) not performed, i.e. bandage, splint will be pointed out to team at this time.
16. The calculated time will be determined by Contest Officials by averaging the working time of all teams participating in the Contest (1 discount per 3 minute overtime or fraction thereof).
17. The accumulation of individual discounts within a procedure shall not exceed the discounts for failure to perform that procedure. (Example AV, CPR, etc.)

A. Ties

In the event of ties in the contest, Scorecard A (First Aid Procedures and Critical Skills) discounts will be the first tie breaker, Scorecard B (AV/CPR) discounts will be the second tie breaker and actual working time, in minutes and seconds, of the team will be the third tie breaker.

B. Miscellaneous

Teams will be notified by posting when they may review their score cards. Within one hour of posting, team members and trainer shall report to a designated location. Once notified, team members and the trainer shall have 20 minutes for reviewing each problem, the judge's skill sheets and scorecards to prepare any protest. All protests shall be in writing and shall state the discount in question, the scorecard involved, and their reference proof in the rule book or Brady book to support their protest or the protest will not be considered. All protests will be considered by the Final Appeals Committee. A decision by the Final Appeals Committee is binding and final. Protest sheets will be furnished to the teams by the Judges for the recording of rules infractions or discounts assessed to teams. Judges shall remain available until released.

MATERIALS LIST

Participants will be required to furnish their own materials. Teams must provide the minimum equipment. Listed below is the minimum equipment required.

- 24 Triangular Bandages
- 6 Adhesive compresses
- 24 Sterile gauze, (4"x4") and/or 4" Compresses
- 6 Roller Bandages
- 2 Blankets
- 1 Scissors, EMT Utility
- 6 Pairs of Examination Gloves
- 2 Mask/face shields or masks and goggles combination meeting blood borne pathogen requirements
- 2 Heat Pack - Simulated
- 2 Cold packs - Simulated
- 2 Oval Eye Pads
- 1 Pen and paper set
- 1 Elevating device
- 1 Recording manikin
- 2 Barrier devices with one-way valve for performing AV/CPR
- 1 White bag (i.e. plastic garbage bag)
- 1 Compliment of splints (may be pre-padded but not assembled)
- 1 Long back board with straps (Aluminum, Wood, etc.)
- 2 Air splints (1 full arm and 1 full leg)
- 1 Packet Sugar/Tube Instant Glucose (for Diabetic Purposes)
- 1 Adhesive Tape
- 1 Burn Sheet, Sterile (40" x 80" minimum)
- 1 Rigid Extrication Collar
- 4 Trauma Dressings (minimum of 10" X 30")
- 1 Eye Shield/Cup
- 1 Pen Light
- 2 Tourniquets
- 2 Towels
- 1 Pillow
- 1 Occlusive Dressing
- 2 Sticks, Wooden Dowels or equivalent
- 1 Watch/Timing Device
- 1 Headset (long spine board)
- 1 500 ml sterile water
- Compliment of Straps for Long Spine Board (buckle straps, spider straps, etc.)

Problems will be designed from the Revised Skill Sheets approved by the Rules Committee.

Manikins will be furnished by teams for performing procedures and critical skills pertaining to all ventilation problems, cardiac arrest problems. Signal boxes on manikins will be covered during the working of the problem. NOTE: Live patients will **not** be used in any CPR or ventilation problems.

Under no circumstances will videotape recordings or photos be introduced as supplementary material for consideration of the appeal.

SCORECARD A DISCOUNTS

1. Violations of general rules not covered on scorecards. ____5 each infraction
2. All life-threatening conditions shall be located and started before patient assessment can begin. (Life threatening conditions will be considered a patient having any one or more of the following conditions: breathing difficulties, no pulse, spinal injury, skull fracture, a sucking chest wound or life threatening bleeding) ____ 20

Patient assessment can begin after all life-threatening conditions have been located and treatment started. Environmental and Medical Emergencies can be treated anytime during the working of the problem after initial assessment.

3. When the team encounters life-threatening bleeding, no work other than controlling bleeding shall be done until bleeding is controlled. Bleeding is controlled when notified by the Judge (judge makes a statement that bleeding is controlled). If treatment has been started and one team member can complete that treatment, the other team member may continue to work. ____10 each infraction
4. During the course of the problem, teams may encounter a card, envelope or label stating various conditions. Upon completion of treatment of these conditions, resume patient assessment at the point where team left off. ____5 each infraction
5. Patient cannot talk, direct, or assist unless stated in the problem. (Reactionary or unintentional movements by the patient should not be discounted) ____ 5 each infraction
6. The bystander must be shown the correct method of support. ____ 2

The bystander must be shown the correct method of support and maintaining the open airway by a team member or members any time during the working of the problem, but before taking support.

7. Handling of a patient by a team or team member in such a manner that could compromise condition of the patient. Examples: Mishandling extremities, etc.) ____5 each infraction
8. All injuries and/or conditions shall be treated (example: wound, fracture, frostbite). ____20 each infraction

9. Preassembly of material. ____5
10. Failure to perform a required critical skill. Each CRITICAL SKILL shall be performed as identified on the skill sheets. ____2 each infraction
11. During patient assessment, failure to verbally state the location physically examined and each condition found. ____1 each infraction
12. Working out of order (assessment, procedure, critical skill). ____2

Low Priority

Teams will systematically conduct the patient assessment according to procedures of the patient assessment skill sheet. Each area of the body shall be examined in its entirety prior to treating injuries in that area (except taking support). All injuries must be treated on the area being examined prior to moving to the next area to be examined. The sling for fractured ribs may be applied after upper extremity has been surveyed/treated. If treatment has been started and can be completed by one team member (except injuries requiring a backboard), the other team member may continue the examination to the next area and begin treatment. (Systemically, legs are treated before the arms.

High Priority

Teams will systematically conduct initial assessment, treating all life-threatening injuries/conditions. When one or more of the conditions listed in rule 24 is encountered the team will perform a rapid patient assessment according to the patient assessment skill sheet. To perform a rapid patient assessment, teams will examine each area of the body in its entirety, verbalizing critical skills and injuries/conditions found. No treatment is required for non life-threatening conditions/injuries found during the rapid patient assessment. Patient is then prepared for transport and/or transported as required by written problem. To prepare for transportation, a team will be required to properly place and secure a patient on a backboard as outlined in the skill sheets, cover with a blanket and lift patient from the floor. After the patient has been lifted from the floor, the team will verbalize - "transporting patient".

13. Failure to follow written instructions. ____5

14. Teams shall not pad around the head and neck of the patient, for a suspected spinal injury, before the patient is placed onto the backboard. ____ 1
15. Teams shall tuck the tails of bandages. ____1 (discount only one (1) point when three or more tails of bandages are not tucked)
16. All material (except manikin) shall be placed behind baseline prior to stopping the timing device. After completing the problem the work area shall be cleaned of ALL material (except manikin), including the infectious waste, which shall be placed in a white trash bag provided by the team. When all materials have been placed behind baseline, a team member shall stop the timing device. The judges and First Aid team will verify the working time upon completion of the problem.
____1
17. Protective equipment must be donned prior to patient contact (gloves, masks, and eye protection - eyeglasses are acceptable). Only BSI may be donned prior to starting the timing device. ____5 each infraction
18. Gloves shall be changed if there would be contamination because of a glove tear or due to other contamination. For contest purposes only, gloves must be changed if there would be provider contamination because of a glove tear. ____2 each infraction (New glove may be placed over original glove)
19. The broken-back board splint may be preassembled and padded. Other splints may be prepped but not assembled. (Cravat bandages cannot be preassembled on the back board, except for tying padding.) ____5
20. Failure to take support of a fracture or dislocation (not supporting fracture or dislocation). ____10
 - Support of Extremities – Above and below the fracture or dislocation
 - Support of Hip – Both sides of the fracture or dislocation
 - Support for spinal injury – Stabilization of neck/Modified Jaw Trust
 - Support for skull fracture – Stabilization of neck/Modified Jaw Trust
 - No support for fractured ribs,
 - No support of fractures/dislocations of nose, jaw, fingers, and toes

21. Support of fractures and/or dislocations shall not be broken or released. ____5

When changing support, if support is broken, this discount applies. Change of support can be done as many times as the team desires provided the support is not broken.

Support for upper extremity fractures/dislocations shall be maintained until the sling is completed. Discount if support of fracture and/or dislocation is released by support person before sling is completed.

Sling and swath not required with air splints.

22. Fractures/dislocations shall be supported prior to bandaging injuries. Once the extremity has been assessed, fractures/dislocations must be supported prior to bandaging injuries on the extremity. ____5

During initial and patient assessment, teams must physically support/stabilize fractures and dislocations that require support as they are found. When the fracture/dislocation is on an extremity and support has been taken, the team must complete the examination on the extremity treating other injuries prior to splinting the fracture/dislocation.

23. Not applying sling for upper extremity wound. ____1

Triangular slings are required for all wounds of upper extremities, including shoulder and armpit wounds. Slings will not be required for upper extremity burns/deep cold injuries. However, if a burn/deep cold injury and wound and/or fracture/dislocation are present on the same upper extremity, a sling shall be applied. Slings will be applied on upper extremities with injuries after all injuries of the extremity have been treated. Slings are not required when the extremity is supported/secured by an air splint or full arm splint.

24. Failure to determine high priority patients. ____10

A high priority patient shall be transported immediately. This presents a load and go situation. High priority conditions are unresponsiveness, breathing difficulties, life threatening bleeding or severe shock, chest pain, no pulse, spinal injury, skull fracture, a sucking chest wound and any severe abdominal, pelvic or spinal pain. Any one or more of the above conditions must be clearly stated in the reading of the problem or written information presented to the team during the working of the problem.

25. Failure to start timing device. ____ 2 discounts

INTERPRETATIONS OF SCORECARD B
ARTIFICIAL VENTILATION/CARDIOPULMONARY RESUSCITATION

1. Failure to determine unresponsiveness (according to Critical Skill Sheet). ____1
2. Failure to call for help. ____1
3. Failure to open airway. ____1
4. Failure to use proper maneuver to open airway (using head-tilt/chin-lift maneuver when jaw-thrust should be used, vice versa). ____1
5. Failure to assess breathlessness within 10 seconds. ____1
6. Failure to use one-way valve barrier device when ventilating manikin. ____1
7. Failure to give initial 2 breaths. ____1
 - a. Volume shall be at least .8 liters (through .7 liter line on new manikins). Over inflation shall not be discounted. ____1
8. Failure to use mouth-to-nose ventilation when required. ____1
9. Failure to keep body and head in line, if spinal injury exists. ____1
10. Failure to use tongue jaw lift, cross-finger technique, or finger sweep when required. ____1
11. Failure to reposition head when airway obstruction is suspected. ____1
12. Failure to give abdominal thrusts or chest thrusts when required. ____1
13. Failure to check pulse prior to giving compressions. ____1
14. Failure to assess pulse for 5-10 seconds. ____1
15. Failure to correctly locate the carotid pulse. ____1
16. Failure to verbalize absence of pulse. ____1

Cardiopulmonary Resuscitation

1. Failure to give AV/CPR when required. ____20
(Maximum of 3 sets AV/CPR or combination thereof)
2. Failure to locate landmark for giving compressions. ____1
3. Failure to make parallel axis with heels of hands. ____1
4. Allowing fingers to rest on chest. ____1
5. Compressions. Discounts shall apply to each set.
 - a. Timing. 30 compressions shall be delivered within 23 seconds. ____1
 - b. Depth. Compression depth shall be between the two lines for 60-80 pounds pressure. ____1
 - c. Number required. A total of 30 compressions shall be made each cycle. ____1
 - d. Release of upstroke. The release line shall be straight. ____1
 - e. Rate. Compressions shall be made at the rate of 100 per minute. ____1
6. Failure to maintain hand contact with manikin when releasing pressure during compressions. ____1 (This does not apply between cycles).
7. Failure to give 2 breaths between each cycle of compressions. ____1
 - a. Timing (not completing breaths and returning to compressions in 4-7 seconds (This will be measured from the end of last down stroke to the start of the first down stroke of the next cycle.) ____1
 - b. Volume shall be at least .8 liters (through .7 liter line on new manikins). Over inflation shall not be discounted. ____1
8. Failure to give 5 cycles of 30 compressions and 2 breaths for each set of CPR (point of first down stroke to peak of last breath). (A cycle is 30 compressions and two (2) ventilations. A set is 5 cycles.) ____1
9. Failure to assess pulse within 10 seconds after each set of CPR. ____1 (one discount per set)

10. Failure to give five (5) abdominal thrusts when airway obstruction is suspected. ____1
11. Failure to perform CPR as stated in the problem. Too many or too few compressions can be detrimental to patient. ____1
12. Failure for the number of Rescuer/Rescuers to perform CPR as stated in the problem. Team performing One-Person CPR when Two-Person CPR is required and vice versa. ____ 3 (When problem states "Two-Rescuer CPR", two people are required to perform CPR as listed in Two-Rescuer CPR skill sheets.)
13. Failure to begin with compressions after pulse check is completed or when changing rescuers. ____1
14. Failure of rescuers to change positions in 5 seconds or less when performing two-person CPR. ____1
15. Failure of rescuer to state that patient has a pulse when CPR is completed. ____1

Artificial Ventilation

1. Failure to give artificial ventilation. _____ 20
(Maximum of 3 sets AV/CPR or combination thereof)
2. Failure to give 10-12 breaths in each 58-62-second period. ____1
(1 minute of AV = 1 set)
3. Failure to provide a breath volume of at least .8 liters (through .7 liter line on new manikins). Over inflation shall not be discounted. _____ 1
4. Failure of rescuer to check for return of breathing and pulse when artificial ventilation is completed. ____1
5. Failure of rescuer to state that patient is breathing and has a pulse when artificial ventilation is completed. ____1

INITIAL ASSESSMENT

PROCEDURES		CRITICAL SKILL
1. SCENE SIZE UP	<input type="checkbox"/> <input type="checkbox"/>	*A. Observe area to ensure safety *B. Call for help
2. MECHANISM OF INJURY	<input type="checkbox"/> <input type="checkbox"/>	*A. Determine causes of injury, if possible *B. Ask patient (if conscious) what happened
3. INITIAL ASSESSMENT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Verbalize general impression of the patient(s) *B. Determine responsiveness/level of consciousness (AVPU) Alert, Verbal, Painful, Unresponsive *C. Determine chief complaint/apparent life threats
4. ASSESS AIRWAY AND BREATHING	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Correctly execute head-tilt/chin-lift or jaw thrust maneuver, depending on the presence of cervical spine (neck) injuries B. Look, listen, and feel for breathing (3-5 seconds) C. If present, treat sucking chest wound
5. ASSESS FOR IMMEDIATE LIFE THREATENING CONDITIONS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Check for presence of a carotid pulse (5-10 seconds) B. If present, control life threatening bleeding C. Start treatment for all other life threatening injuries/conditions (reference Rule 2).
6. DETERMINE PRIORITY OF PATIENT	<input type="checkbox"/> <input type="checkbox"/>	*A. Teams must make statement to judge, identifying whether patient is low priority or high priority load and go. *B. Teams must make statement to judge, "Removing clothing, exposing and cleaning wound surface(s)"

HIGH PRIORITY: Rapid Patient Assessment treating all life threats Load and Go.
If the treatment interrupts the rapid trauma assessment, the **assessment** will be completed at the end of the **treatment**.

LOW PRIORITY: Detailed Patient Assessment treating all injuries and conditions and prepare for transport.

14

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

PATIENT ASSESSMENT

PROCEDURES		CRITICAL SKILL	
1. HEAD	<div><input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/></div>	<div>*A. Check head for DOTS: Deformities, Open wounds, Tenderness and Swelling</div> <div>*B. Check and touch the scalp</div> <div>*C. Check the face</div> <div>*D. Check the ears for bleeding or clear fluids</div> <div>*E. Check the eyes for any discoloration, unequal pupils, reaction to light, foreign objects and bleeding</div> <div>*F. Check the nose for any bleeding or drainage</div> <div>*G. Check the mouth for loose or broken teeth, foreign objects, swelling or injury of tongue, unusual breath odor and discoloration</div>	
2. NECK	<div><input type="checkbox"/> <input type="checkbox"/></div>	<div>*A. Check the neck for DOTS</div> <div>*B. Inspect for medical ID</div>	
3. CHEST	<div><input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/></div>	<div>*A. Check chest area for DOTS</div> <div>*B. Feel chest for equal breathing movement on both sides</div> <div>*C. Feel chest for inward movement in the rib areas during inhalations</div>	
4. ABDOMEN	<div><input type="checkbox"/></div>	<div>*A. Check abdomen (stomach) for DOTS</div>	
5. PELVIS	<div><input type="checkbox"/> <input type="checkbox"/></div>	<div>*A. Check pelvis for DOTS</div> <div>*B. Inspect pelvis for injury by touch (Verbally state inspection of crotch and buttocks areas)</div>	
6. LEGS	<div><div><div>L</div><div><input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/></div></div><div><div>R</div><div><input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/></div></div></div>	<div>*A. Check each leg for DOTS</div> <div>B. Inspect legs for injury by touch</div> <div>C. Unresponsive: Check legs for paralysis (pinch inner side of leg on calf)</div> <div>*D. Responsive: Check legs for motion; places hand on bottom of each foot and states “Can you push against my hand?”</div> <div>*E. Check for medical ID bracelet</div>	

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

7. ARMS	L <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	R <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<p>*A. Check each arm for DOTS</p> <p>B. Inspect arms for injury by touch</p> <p>C. Unresponsive: Check arms for paralysis (pinch inner side of wrist)</p> <p>*D. Responsive: Check arms for motion (in a conscious patient; team places fingers in each hand of patient and states "Can you squeeze my fingers?")</p> <p>*E. Check for medical ID bracelet</p>
8. BACK SURFACES	<input type="checkbox"/>		<p>*A. Check back for DOTS</p>

DOTS: Deformities, Open Wounds, Tenderness and Swelling

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

ONE-PERSON CPR (MANIKIN ONLY)

PROCEDURES		CRITICAL SKILL
1. ESTABLISH UNRESPONSIVENESS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Tap or gently shake shoulders *B. "Are you OK?" C. Determine unconsciousness without compromising possible cervical spine (neck) injury *D. "Call for help"
2. ESTABLISH AIRWAY	<input type="checkbox"/> <input type="checkbox"/>	A. Kneel at the patient's side near the head B. Correctly execute head-tilt/chin-lift or jaw thrust maneuver depending on the presence of cervical spine injuries
3. MONITOR PATIENT FOR BREATHING	<input type="checkbox"/>	A. Look, listen, and feel for breathing (within 10 seconds)
4. VENTILATION PATIENT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place barrier device (pocket mask/shield with one way valve) on manikin B. Give 2 breaths 1 second each C. Each breath - minimum of .8 liters (through .7 liter line on new manikins)
5. PULSE CHECK	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Correctly locate the carotid pulse - on the side of the rescuer, locate the patient's windpipe with your index and middle fingers and slide your fingers in the groove between the windpipe and muscle in the neck B. Check for presence of carotid pulse for 5 to 10 seconds *C. Absence of pulse
6. POSITION FOR COMPRESSIONS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Locate the compression point on the breastbone between the nipples B. Place the heel of one hand on the compression point and the other hand on top of the first so hands are parallel C. Do not intentionally rest fingers on the chest D. Keep heel of your hand on chest during and between compressions

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

7. DELIVER CARDIAC COMPRESSION	<input type="checkbox"/> A. Give 30 compressions <input type="checkbox"/> B. Compressions are at the rate of 100 per minute (30 compressions delivered within 23 seconds) <input type="checkbox"/> C. Down stroke for compression must be on or between compression lines <input type="checkbox"/> D. Return to baseline on upstroke of compression
7. VENTILATIONS BETWEEN COMPRESSIONS	<input type="checkbox"/> A. Give 2 breaths 1 second each <input type="checkbox"/> B. Each breath - minimum of .8 (through .7 liter line on new manikins) <input type="checkbox"/> C. Complete breaths and return to compressions in 4-7 seconds (This will be measured from the end of last down stroke to the start of the first down stroke of the next cycle.)
9. CONTINUE CPR FOR TIME STATED IN PROBLEM	<input type="checkbox"/> A. Provide 5 cycles of 30 chest compressions and 2 rescue breaths <input type="checkbox"/> B. To check for pulse, stop chest compressions for 5-10 seconds after the first set of CPR <input type="checkbox"/> C. Rescuer opens airway and looks, listens, and feels for adequate breathing or coughing <input type="checkbox"/> D. Rescuer checks for a carotid pulse <input type="checkbox"/> E. If no signs of circulation are detected, continue chest compressions and breaths and check for signs of circulation after each set <input type="checkbox"/> F. A maximum of 10 seconds will be allowed to complete ventilations and required pulse checks between sets (this will be measured from the end of the last down stroke to the start of the first down stroke of the next cycle)
9. CHECK FOR RETURN OF PULSE	<input type="checkbox"/> A. After providing required CPR (outlined in problem), check for return of pulse (within 10 seconds) <input type="checkbox"/> *B. "Patient has a pulse."

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

TWO-RESCUER CPR (NO SPINAL INJURY - MANIKIN ONLY)

PROCEDURES		CRITICAL SKILL
1. RESCUER 1 - ESTABLISH UNRESPONSIVENESS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Tap or gently shake shoulders *B. "Are you OK?" C. Determine unconsciousness without compromising cervical spine (neck) injury *D. "Call for help"
2. RESCUER 1 - ESTABLISH AIRWAY	<input type="checkbox"/> <input type="checkbox"/>	A. Kneel at the patient's side near the head B. Correctly execute head-tilt/chin-lift maneuver
3. RESCUER 1 - MONITOR PATIENT FOR BREATHING		A. Look, listen, and feel for breathing (within 10 seconds)
4. RESCUER 1 - VENTILATE PATIENT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place barrier device (pocket mask/shield with one-way valve) on manikin B. Give 2 breaths 1 second each C. Each breath - minimum of .8 (through .7 liter line on new manikins)
5. RESCUER 1 - CHECK FOR CAROTID PULSE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Correctly locate the carotid pulse - on the side of the rescuer, locate the patient's windpipe with your index and middle fingers and slide your fingers in the groove between the windpipe and the muscle in the neck B. Check for presence of carotid pulse for 5 to 10 seconds *C. Absence of pulse
6. RESCUER 2 - POSITION FOR COMPRESSIONS	<input type="checkbox"/> <input type="checkbox"/>	A. Locate the compression point on the breastbone between the nipples B. Place the heel of one hand on the compression point and the other hand on top of the first so hands are parallel. Do not intentionally rest fingers on the chest. Keep heel of your hand on chest during and between compressions.

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

7. RESCUER 2 - DELIVER CARDIAC COMPRESSION	<input type="checkbox"/> A. Give 30 compressions <input type="checkbox"/> B. Compressions are at the rate of 100 per minute (30 compressions delivered within 23 seconds) <input type="checkbox"/> C. Down stroke for compression must be on or between compression lines <input type="checkbox"/> D. Return to baseline on upstroke of compression
8. RESCUER 1 - VENTILATIONS BETWEEN COMPRESSIONS	<input type="checkbox"/> A. Give 2 breaths 1 second each <input type="checkbox"/> B. Each breath - minimum of .8 (through .7 liter line on new manikins) <input type="checkbox"/> C. Complete breaths and return to compressions in 4-7 seconds (This will be measured from the end of last down stroke to the start of the first down stroke of the next cycle.)
9. CONTINUE CPR FOR TIME STATED IN PROBLEM	<input type="checkbox"/> A. Provide 5 cycles of 30 chest compressions and 2 rescue breaths <input type="checkbox"/> B. To check for pulse, stop chest compressions for 5-10 seconds after the first set of CPR <input type="checkbox"/> C. Rescuer at patient's head maintains airway and looks, listens, and feels for adequate breathing or coughing <input type="checkbox"/> D. The rescuer at the patient's head shall feel for a carotid pulse <input type="checkbox"/> E. If no signs of circulation are detected, continue chest compressions and breaths and check for signs of circulation after each set <input type="checkbox"/> F. A maximum of 10 seconds will be allowed to complete ventilations and required pulse checks between sets (this will be measured from the end of the last down stroke to the start of the first down stroke of the next cycle)
10. CHANGING RESCUERS	<input type="checkbox"/> A. Change of rescuers shall be made in 5 seconds or less and will be completed as outlined in the problem. Team must switch every 5 cycles in less than 5 seconds.
11. CHECK FOR RETURN OF PULSE	<input type="checkbox"/> A. After providing required CPR (outlined in problem), check for return of pulse (within 10 seconds) <input type="checkbox"/> *B "Patient has a pulse."

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

TWO-RESCUER CPR (WITH SPINAL INJURY - MANIKIN ONLY)

PROCEDURES		CRITICAL SKILL
1. RESCUER 1 - ESTABLISH UNRESPONSIVENESS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Tap or gently shake shoulders *B. "Are you OK?" C. Determine unconsciousness without compromising cervical spine (neck) injury *D. "Call for help"
2. RESCUER 2 - ESTABLISH AIRWAY	<input type="checkbox"/> <input type="checkbox"/>	A. Kneel at the patient's head B. Correctly execute jaw thrust maneuver
3. RESCUER 1 - MONITOR PATIENT FOR BREATHING	<input type="checkbox"/>	A. Look, listen, and feel for breathing (within 10 seconds)
4. RESCUER 2 - VENTILATE PATIENT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Rescuer 1 should place barrier device (pocket mask/shield with one-way valve) on manikin. (OPTION 1: When spinal injury is present, Rescuer No. 2 can hold barrier device on manikin after Rescuer No. 1 correctly places the device over the mouth and nose.) (OPTION 2: Rescuer 1 can place the device on the manikin each time patient is ventilated) B. Rescuer 2 gives 2 breaths 1 second each C. Each breath - minimum of .8 (through .7 liter line on new manikins)
5. RESCUER 1 - CHECK FOR CAROTID PULSE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Correctly locate the carotid pulse - on the side of the rescuer, locate the patient's windpipe with your index and middle fingers and slide your fingers in the groove between the windpipe and the muscle in the neck B. Check for presence of carotid pulse for 5 to 10 seconds *C. Absence of pulse
6. RESCUER 1 - POSITION FOR COMPRESSIONS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Locate the compression point on the breastbone between the nipples B. Place the heel of one hand on sternum the compression point and the other hand on top of the first so hands are parallel C. Do not rest fingers on the chest Keep heel of your hand on chest during and between compressions

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

7. RESCUER 1 - DELIVER CARDIAC COMPRESSION	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Give 30 compressions B. Compressions are at the rate of 100 per minute (30 compressions delivered within 23 seconds) C. Down stroke for compression must be on or between compression lines D. Return to baseline on upstroke of compression
8. RESCUER 2 - VENTILATIONS BETWEEN COMPRESSIONS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Give 2 breaths 1 second each B. Each breath - minimum of .8 (through .7 liter line on new manikins) C. Complete breaths and return to compressions in 4-7 seconds (This will be measured from the end of last down stroke to the start of the first down stroke of the next cycle.)
9. CONTINUE CPR FOR TIME STATED IN PROBLEM	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Provide 5 cycles of 30 chest compressions and 2 rescue breaths B. To check pulse, stop chest compressions for 10 seconds after the first set of CPR C. Rescuer at patient's head maintains airway and looks, listens, and feels for adequate breathing or coughing D. The rescuer giving compressions shall feel for a carotid pulse E. If no signs of circulation are detected, continue chest compressions and breaths and check for signs of circulation after each set F. A maximum of 10 seconds will be allowed to complete ventilations and required pulse checks between sets (this will be measured from the end of the last down stroke to the start of the first down stroke of the next cycle)
10. CHANGING RESCUERS	<input type="checkbox"/>	A. Change of rescuers shall be made in 5 seconds or less and will be completed as outlined in problem. Team must switch every 5 cycles in less than 5 seconds.
11. CHECK FOR RETURN OF PULSE	<input type="checkbox"/> <input type="checkbox"/>	A. A final pulse check (3-5 seconds) will be required at the end of the last set of CPR *B. "Patient has a pulse."

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

MOUTH-TO-MASK RESUSCITATION

PROCEDURES		CRITICAL SKILL
1. ESTABLISH UNRESPONSIVENESS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Tap or gently shake shoulders *B. "Are you OK?" C. Determine unconsciousness without compromising C-spine injury *D. "Call for help"
2. ESTABLISH AIRWAY	<input type="checkbox"/>	A. Correctly execute head-tilt/chin-lift or jaw thrust maneuver depending on the presence of cervical spine (neck) injuries
3. MONITOR PATIENT FOR BREATHING	<input type="checkbox"/>	A. Look, listen, and feel for breathing (within 10 seconds)
4. VENTILATE PATIENT	<input type="checkbox"/> <input type="checkbox"/>	A. Place barrier device (pocket mask/shield with one-way valve) on manikin B. Ventilate patient 2 times at 1 second intervals each - minimum of .8 (through .7 liter line on new manikins)
5. CHECK FOR CAROTID PULSE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Correctly locate the carotid pulse (on the side of the rescuer) B. Check for presence of carotid pulse within 10 seconds *C. Presence of pulse
6. VENTILATE PATIENT	<input type="checkbox"/> <input type="checkbox"/>	A. Place barrier device (pocket mask/shield with one-way valve on manikin B. Ventilate patient 10 to 12 times per minute. Each ventilation will be provided at a minimum of .8 (through .7 liter line on new manikins)
7. CHECK FOR RETURN OF BREATHING AND PULSE	<input type="checkbox"/> <input type="checkbox"/>	A. After providing the required number of breaths (outlined in problem), check for return of breathing and carotid pulse within 10 seconds *B. "Patient is breathing and has a pulse"

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

AIRWAY OBSTRUCTION (MANIKIN ONLY) (VICTIM LOSES CONSCIOUSNESS)

PROCEDURES		CRITICAL SKILL
1. POSITIONING VICTIM AFTER LOSS OF CONSCIOUSNESS	<input type="checkbox"/> <input type="checkbox"/>	A. Place victim in supine position (on their back) *B. "Call for help"
2. PERFORM FINGER SWEEPS	<input type="checkbox"/> <input type="checkbox"/>	A. Follow with opening mouth and finger sweep (Open mouth, grasping tongue and lower jaw with thumb and fingers, insert index finger of other hand down along inside cheek and deeply into throat in a hooking action) B. Grasp and remove foreign object if dislodged
3. OPEN AIRWAY	<input type="checkbox"/>	A. Correctly execute head-tilt/chin-lift or jaw thrust maneuver depending on the presence of cervical spine (neck) injuries
4. ATTEMPT VENTILATIONS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place barrier device on manikin B. Seal mouth and nose C. Attempt to give slow full breath D. If successful, deliver second slow full breath and check pulse E. If unsuccessful, continue sequence
5. DELIVER ABDOMINAL THRUST	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Straddle the victim's thighs B. Place heel of one hand against the victim's abdomen, in midline slightly above the navel and well below Xiphoid (point where ribs attach to the lower part of the breastbone) C. Place second hand on top of the first D. Deliver quick upward thrust E. Each thrust administered with intent of relieving obstruction F. Deliver 5 thrusts
6. PERFORM FINGER SWEEP	<input type="checkbox"/> <input type="checkbox"/>	A. Follow with opening mouth and finger sweep (open mouth, grasping tongue and lower jaw with thumb and fingers, insert index finger of other hand down along inside cheek and deeply into throat in a hooking action) B. Grasp and remove foreign object if dislodged

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

7. ATTEMPT VENTILATIONS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place barrier device on manikin B. Correctly make effort to administer slow full breath C. If successful, deliver second slow full breath and check pulse D. If unsuccessful, continue sequence
8. REPEAT SEQUENCE UNTIL SUCCESSFUL	<input type="checkbox"/>	A. Demonstrate ability to rapidly repeat sequence 1-7
9. ESTABLISH AIRWAY	<input type="checkbox"/> <input type="checkbox"/>	A. Correctly execute head-tilt/chin-lift or jaw thrust maneuver depending on the presence of cervical spine (neck) injuries B. Look, listen, and feel for the presence of breathing (within 10 seconds)
10. RE-EVALUATION OF PATIENT	<input type="checkbox"/> <input type="checkbox"/>	*A. Determine the need to repeat the sequence B. Cease at the direction of the judge

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

AIRWAY OBSTRUCTION (UNCONSCIOUS VICTIM - WITNESSED)

PROCEDURES		CRITICAL SKILL
1. INITIALLY ASSESS LEVEL OF CONSCIOUSNESS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Tap or gently shake shoulders *B. "Are you OK?" C. Determine unconsciousness without compromising C-spine injury *D. "Call for help"
2. ESTABLISH AIRWAY	<input type="checkbox"/>	A. Correctly execute head-tilt/chin-lift or jaw thrust maneuver depending on the presence of cervical spine (neck) injuries
3. MONITOR PATIENT FOR BREATHING	<input type="checkbox"/> <input type="checkbox"/>	A. Look, listen, feel for presence of breathing (within 10 seconds) B. Maintain open airway
4. ATTEMPT VENTILATION	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place barrier device on manikin B. Seal mouth and nose C. Attempt to give slow breath *D. Identify if there is an obstruction
5. CHECK POSITIONING	<input type="checkbox"/> <input type="checkbox"/>	A. Re-establish airway using correct method and procedure B. Identify continued presence of the obstruction
6. DELIVER ABDOMINAL THRUST	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Kneel astride the victim's thighs B. Place heel of one hand against the victim's abdomen, in midline slightly above the navel and well below Xiphoid (point where ribs attach to the lower part of the breastbone) C. Place second hand on top of the first D. Deliver quick upward thrust E. Each thrust administered with intention of relieving obstruction F. Deliver 5 thrusts

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

7. DELIVER CHEST THRUST (ONLY USED IN ADVANCED PREGNANCY OR MARKED OBESE VICTIM)	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Kneel at side of victim B. Place heel of hand on breastbone between the nipples (CPR position) C. Place heel of other hand directly on top of the first hand D. Deliver slow distinct downward thrust, with each thrust administered with intent of relieving obstruction E. Deliver 5 thrusts
8. CHECK FOR FOREIGN BODY	<input type="checkbox"/> <input type="checkbox"/>	A. Follow with opening mouth and finger sweep (Open mouth, grasping tongue and lower jaw with thumb and fingers, insert index finger of other hand down along inside cheek and deeply into throat in a hooking action) *B. Grasp and remove foreign object if dislodged
9. ESTABLISH AIRWAY	<input type="checkbox"/>	A. Correctly perform head-tilt/chin-lift or jaw thrust maneuver depending on cervical spine (neck) injuries
10. ATTEMPT VENTILATION	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Correctly make effort to administer slow full breath B. Administer second slow full breath, if first successful and check pulse C. If unsuccessful repeat sequence of thrust, finger sweep and attempt to ventilate

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

SUCKING CHEST WOUND

PROCEDURES		CRITICAL SKILL
1. EXPOSE WOUND	<input type="checkbox"/>	*A. Expose entire wound
2. SEAL WOUND AND CONTROL BLEEDING	<input type="checkbox"/> <input type="checkbox"/>	*A. Place occlusive dressing over wound (If occlusive dressing is not available use gloved hand) B. Apply direct pressure as needed to stop the bleeding
3. APPLY AN OCCLUSIVE DRESSING	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Keep patient calm and quiet *B. Explain to the patient what you are doing *C. Ensure dressing is large enough not to be sucked into the wound (two inches beyond edges of wound) D. Affix dressing with tape *E. Seal on three sides *F. Monitor patient closely for increasing difficulty breathing *G. Transport as soon as possible H. Keep patient positioned on the injured side unless other injuries prohibit *I. Reassess wound to ensure bleeding control *J. Assess level of consciousness(AVPU), respiratory status and patient response

NOTE: Alert, verbal, painful, unresponsive (AVPU)

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

LIFE-THREATENING BLEEDING

PROCEDURES	CRITICAL SKILL
1. DIRECT PRESSURE AND ELEVATION	<input type="checkbox"/> *A. Expose the wound. <input type="checkbox"/> *B. Clear the wound surface <input type="checkbox"/> C. Apply direct pressure with a gloved hand <input type="checkbox"/> D. Apply a dressing to wound (cover entire wound) and continue to apply direct pressure <input type="checkbox"/> E. Elevate the extremity except when spinal injury exists <input type="checkbox"/> *F. Bleeding has been controlled <input type="checkbox"/> G. If controlled, bandage dressing in place
2. IF NOTIFIED THAT BLEEDING IS NOT CONTROLLED, PRESSURE POINTS SHALL BE UTILIZED	<input type="checkbox"/> *A. Apply pressure to appropriate pressure point and notify judge verbally that bleeding is controlled (Apply pressure to blood vessels leading to area – in arm, press just below armpit; in leg, press against groin where thigh and trunk join.) <input type="checkbox"/> B. If controlled, bandage dressing in place
3. IF NOTIFIED THAT BLEEDING IS NOT CONTROLLED, APPLY TOURIQUET	<input type="checkbox"/> A. Apply as per tourniquet skill sheet

External Bleeding

To Control: 1st: direct pressure
 2nd: elevation & direct pressure
 3rd: pressure point
 Last Resort: Tourniquet

Internal Bleeding

- *1. Monitor breathing and pulse
2. Keep patient still
- *3. Loosen restrictive clothing
4. Be alert if patient vomits
5. Nothing by mouth
6. Report possibility of internal bleeding as soon as EMS personnel arrive on scene

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

TOURNIQUET

PROCEDURES		CRITICAL SKILL
1. DETERMINE NEED FOR USING TOURNIQUET	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<p>If these conditions are met, a tourniquet may be the only alternative:</p> <p>A. Direct pressure has not been successful in stopping bleeding</p> <p>B. Elevation of wound above heart has not been successful in stopping of bleeding</p> <p>C. Compression of pressure point has not been successful in stopping of bleeding.</p>
2. SELECT APPROPRIATE MATERIALS	<input type="checkbox"/>	<p>A. Select a band that will be between 3-4 inches in width and can be wrapped six or eight layers deep for improvised tourniquet or select factory tourniquet.</p>
3. APPLY TOURNIQUET	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<p>Factory Tourniquet</p> <p>A. Wrap band around the extremity proximal to the wound (one inch above but not on a joint)</p> <p>Improvised Tourniquet</p> <p>B. Apply a bandage around the extremity proximal to the wound (one inch above but not on a joint) and tie a half knot in the bandage</p> <p>C. Place a stick or pencil on top of the knot and tie the ends of the bandage over the stick in a square knot</p> <p>D. Twist the stick until the bleeding is controlled, secure the stick in position</p>
4. APPLY PRESSURE	<input type="checkbox"/> <input type="checkbox"/>	<p>A. Do not cover the tourniquet</p> <p>*B. Notify other medical personnel caring for the patient</p>
5. MARK PATIENT APPROPRIATELY	<input type="checkbox"/>	<p>A. Mark a piece of tape on the patient's forehead "TQ" and time applied</p>
6. REASSESS	<input type="checkbox"/>	<p>*A. Assess level of consciousness, respiratory status, and patient response</p>

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

DRESSINGS AND BANDAGING - OPEN WOUNDS

PROCEDURES	CRITICAL SKILL
1. EMERGENCY CARE FOR AN OPEN WOUND	<input type="checkbox"/> *A. Expose wound <input type="checkbox"/> *B. Clear wound surface <input type="checkbox"/> *C. Control bleeding <input type="checkbox"/> *D. Prevent further contamination <input type="checkbox"/> E. Bandage dressing in place after bleeding has been controlled <input type="checkbox"/> F. Keep patient lying still
2. APPLY DRESSING	<input type="checkbox"/> A. Use sterile dressing <input type="checkbox"/> B. Cover entire wound <input type="checkbox"/> C. Control bleeding <input type="checkbox"/> D. Do not remove dressing
3. APPLY BANDAGE	<input type="checkbox"/> A. Do not bandage too tightly <input type="checkbox"/> B. Do not bandage too loosely <input type="checkbox"/> C. Do not leave loose ends <input type="checkbox"/> D. Cover all edges of dressing <input type="checkbox"/> E. Do not cover tips of fingers and toes, unless they are injured <input type="checkbox"/> F. Bandage from the bottom of the limb to the top (distal to proximal) if applicable

Impaled Objects

- *1. Do not remove
2. Expose wound
3. Control bleeding
4. Stabilize with a bulky dressing; criss-cross the layers
5. Tie 4in. wide cravats around to hold in place, or tape in place
- *6. Check for exit wound
7. Immobilize affected area

Impaled Objects in the Cheek

- *1. Examine; inside & outside
2. If end not impaled in mouth – pull it out
3. Position head for drainage: if spinal injury, immobilize 1st and tilt board
4. Dress outside of wound
- *5. Gauze on inside only if patient alert, (Simulate only in contest and state, “I would leave 3-4 inches of gauze outside of mouth.”)

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

Impaled Objects in the Eye

1. Stabilize with 3 inch gauze or folded 4x4
2. Put cup (no Styrofoam) over object and allow cup to rest on roller gauze or 4x4
3. Secure cup with roller gauze (not over top of cup)
- *4. Cover uninjured eye too

Open Neck Wound (Serious or Life Threatening)

- *1. Gloved hand over wound
- *2. Occlusive dressing over wound- 2 inches larger than wound site
3. Gauze dressing over occlusive
4. Place roller gauze beside site and wrap around figure 8 under opposite arm

Abdominal Injury

- *1. Place on back with legs flexed at the knees (for closed or open wounds)

Additional Steps for Open Abdominal Wounds (Serious or Life Threatening)

- *1. Apply moist dressing, then an occlusive dressing
- *2. Cover the occlusive with pads or a towel for warmth
- *3. If an object is impaled in abs, stabilize it and do not flex legs- leave them in the position you found them.

Skull Fractures and Brain Injuries

- *1. Open airway with jaw thrust
2. Apply collar
- *3. Use loose gauze dressing- no direct pressure
- *4. Keep at rest, ask them questions
5. Don't elevate legs (on or off a backboard)
6. After entire body is immobilized- tilt back board, injured side down

Amputations

- *1. Wrap in slightly moistened sterile dressing
2. Place in plastic bag or wrap in plastic
- *3. Keep part cool avoid freezing
- *4. Do not place in water or direct contact with ice
- *5. Transport with patient
6. Label with patients name

NOTE:

Triangular slings are required for all wounds of upper extremities, including shoulder and armpit wounds. Slings will not be required for upper extremity burns. However, if a burn and wound and/or fracture/dislocation are present on the same upper extremity, a sling shall be applied. Slings will be applied on upper extremities with injuries after all injuries of the extremity have been treated. (Page # 9 rule #24)

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

TWO-PERSON LOG ROLL

PROCEDURES		CRITICAL SKILL
1. STABILIZE HEAD	<input type="checkbox"/>	*A. Stabilize the head and neck
2. PREPARING THE PATIENT	<input type="checkbox"/> <input type="checkbox"/>	A. When placing patient on board place board parallel to the patient B. Kneel at the patient's shoulders opposite the board (if used) leaving room to roll the patient toward knees. Raise the patient's arm, if not injured (the one closer to the rescuer) above the patient's head
3. PREPARING THE RESCUER	<input type="checkbox"/> <input type="checkbox"/>	A. Grasp the patient at the shoulder and pelvis area B. Give instructions to bystander, if used to support
4. ROLLING THE PATIENT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. While stabilizing the head, roll the patient toward the rescuer by pulling steadily and evenly at the shoulder and pelvis areas B. The head and neck should remain on the same plane as the torso C. Maintain stability by holding patient with one hand and placing board (if used) with other D. Roll the body as a unit onto the board (if used) (board may be slanted or flat) E. Place the arm alongside the body

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

SPLINTING (RIGID) UPPER EXTREMITY FRACTURES AND DISLOCATIONS

PROCEDURES		CRITICAL SKILL
1. CARE FOR FRACTURE	<input type="checkbox"/>	*A. Check for distal circulation, sensation, and motor function ■ Do not attempt to reduce dislocations (if applies)
2. IMMOBILIZING FRACTURE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Selection of appropriate rigid splint of proper length B. Support affected limb and limit movement C. Apply appropriate padded rigid splint against injured extremity D. Place appropriate roller bandage in hand to ensure the position of function E. Secure splint to patient with roller bandage, handkerchiefs, cravats, or cloth strips F. Apply wrap distal to proximal *G. Recheck distal circulation, sensation, and motor function
3. SECURING WITH SLING	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place sling over chest and under arm B. Hold or stabilize arm C. Triangle should extend behind elbow or injured side D. Pull sling around neck and tie on uninjured side E. Pad at the neck (except when C-Collar is present) F. Secure excess material at elbow G. Fingertips should be exposed *H. Check for motion and circulation
4. SECURING SLING WITH SWATHE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Use triangle cravat B. Swathe is tied around chest and injured arm *C. Reassess distal circulation, sensation, and motor function

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

ELBOW (STRAIGHT POSITION)

Follow Procedures No. 1 and No. 2 above

FINGER/FINGERS

Immobilize Fracture

1. Tape injured finger to an adjacent uninjured finger; or
2. Tape injured finger to a tongue depressor, aluminum splint, or pen and pencil
3. Secure with sling and swathe

COLLAR BONE

Support and limit movement of affected area

Follow Procedures No. 1, No. 3 and No. 4 above

SHOULDER BLADE

Support and limit movement of affected area

Follow Procedures No. 1, No. 3 and No. 4 above

NOTE: Do not reposition dislocations

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

SPLINTING (SOFT) UPPER EXTREMITY FRACTURES AND DISLOCATIONS (WRIST AND HAND)

PROCEDURES		CRITICAL SKILL
1. CARE FOR FRACTURE	<input type="checkbox"/>	*A. Check for distal circulation, sensation, and motor function <ul style="list-style-type: none"> ▪ Do not attempt to reduce dislocations (if applies)
2. IMMOBILIZING FRACTURE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Support affected limb and limit movement B. Place two cravats (triangular bandage) under wrist/hand C. Place pillow length wise under wrist/hand, on top of cravats (pillow should extend past fingertips) D. Lower limb, adjust cravats to tie E. Tie cravats distal to proximal
3. SECURING WITH SLING	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place sling over chest and under arm B. Hold or stabilize arm C. Triangle should extend behind elbow or injured side D. Secure excess material at elbow E. Fingertips should be exposed *F. Check for motion and circulation
4. SECURING SLING WITH SWATHE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Use triangle cravat B. Swathe is tied around chest and injured arm *C. Reassess distal circulation, sensation, and motor function

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

SPLINTING (RIGID OR SOFT)

PELVIC GIRDLE, THIGH, KNEE, AND LOWER LEG

PROCEDURE		CRITICAL SKILL	
1. DETERMINE NEED FOR SPLINTING	<div><input type="checkbox"/></div> <div><input type="checkbox"/></div>	<div>*A. Assess for:<div><div>▪ Pain</div><div>▪ Swelling</div><div>▪ Deformity</div></div></div> <div>B. Determine if splinting is warranted</div>	
2. APPLY MANUAL STABILIZATION	<div><input type="checkbox"/></div>	<div>A. Support affected limb and limit movement<div><div>▪ Do not attempt to reduce dislocations</div></div></div>	
3. SELECT APPROPRIATE SPLINT	<div><input type="checkbox"/></div> <div><input type="checkbox"/></div>	<div>A. Select appropriate splinting method depending on position of extremity and materials available</div> <div>B. Select appropriate padding material</div>	
4. PREPARE FOR SPLINTING	<div><input type="checkbox"/></div> <div><input type="checkbox"/></div> <div><input type="checkbox"/></div> <div><input type="checkbox"/></div> <div><input type="checkbox"/></div>	<div>A. Remove or cut away clothing as needed</div> <div>*B. Assess PMS distal to the injury:<div><div>▪ Pulse</div><div>▪ Motor Function</div><div>▪ Sensory Function</div></div></div> <div>C. Cover any open wounds with sterile dressing and bandage</div> <div>D. Measure splint</div> <div>E. Pad around splint for patient comfort</div>	

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

SPLINTING (SOFT) LOWER EXTREMITY FRACTURES AND DISLOCATIONS (ANKLE AND FOOT)

PROCEDURES		CRITICAL SKILL
1. CARE FOR FRACTURE	<input type="checkbox"/>	*A. Check for distal circulation, sensation, and motor function <ul style="list-style-type: none"> ▪ Do not attempt to reduce dislocations (if applies)
2. IMMOBILIZING FRACTURE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Support affected limb and limit movement B. Place three cravats (triangular bandage) under ankle/foot C. Place pillow length wise under ankle/foot, on top of cravats (pillow should extend 6 inches beyond foot) D. Lower limb, adjust cravats to tie E. Tie cravats distal to proximal F. Elevate with blanket or pillow *G. Reassess distal circulation, sensation, and motor function

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

SPLINTING UPPER EXTREMITY/LOWER EXTREMITY FRACTURES (AIR SPLINT)

PROCEDURES		CRITICAL SKILL
1. CARE FOR FRACTURE	<input type="checkbox"/>	*A. Check for motion and circulation (fingers/toes)
2. IMMOBILIZE FRACTURE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Grasp above and below the injury site B. Maintain support C. Properly apply air splint D. Splint should be relatively free of wrinkles E. Inflate splint to point that slight dent can be made *F. Reassess circulation (fingers/toes)
3. MONITOR AIR-INFLATED SPLINT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Periodically check for increase or decrease in pressure *B. Monitor pressure in splint with finger tip C. Make sure desired pressure is maintained *D. Reassess circulation (fingers/toes)

NOTE: Air splints may not be used with open (protruding bones) fractures.
Air splints may only be used on the lower part of the extremities (from below the elbow on the arm and below the knee to the leg).

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

SPLINTING - FLAIL CHEST

PROCEDURES	CRITICAL SKILL
1. DETERMINE NEED FOR SPLINTING	*A. Assess for: <ul style="list-style-type: none"> • Pain • Swelling • Deformity *B. Determine if splinting is warranted
2. SELECT APPROPRIATE SPLINTING MATERIAL	A. Choose a pillow, blanket, trauma dressing, or other appropriate splinting material
3. PREPARE FOR SPLINTING	*A. Remove or cut away clothing as needed. B. Cover any open wounds with sterile dressing and bandage
4. APPLY SPLINT	A. Affix splint to chest with adhesive tape or roller bandage B. Immobilize the site of injury C. Use caution when taping splint to chest circumferentially D. Ensure sufficient chest expansion
5. REASSESS	*A. Assess patient response and level of comfort
6. ASSIST VENTILATIONS	A. Assist with ventilation as needed

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

ONE RESCUER BLANKET DRAG

PROCEDURES		CRITICAL SKILL
1. VICTIM SUPINE ON GROUND	<input type="checkbox"/> <input type="checkbox"/>	A. Properly prepare blanket for use in blanket drag B. Spread blanket along side patient with approximately one half the width gathered lengthwise into pleats
2. POSITION PATIENT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Properly roll victim on one side B. Take patients arm on side of body opposite to blanket and extend arm over head C. Support head and neck roll patient on side away from Blanket
3. PLACE PATIENT ON BLANKET	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Properly position on blanket B. Hold patient on side while pleated portion of blanket is pulled in close to victim's back C. Roll patient onto blanket, extend opposite arm and roll onto opposite side D. Smooth out pleats and roll patient onto back E. Snugly wrap patient in blanket with arms at sides
4. PREPARE TO DRAG PATIENT	<input type="checkbox"/> <input type="checkbox"/>	A. Proper blanket drag of patient B. Grasp portion of blanket beneath victim's head and drag victim to safety

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

TWO RESCUER EXTREMITY GROUND LIFT

PROCEDURES		CRITICAL SKILL
1. POSITIONING	<input type="checkbox"/> <input type="checkbox"/>	A. Rescuer 1 – Kneel at the head of the patient and place one hand under each of the shoulders B. Rescuer 2 – Kneel by the patients knees and grasp the patient’s wrist
2. RAISING PATIENT TO A SITTING POSITION	<input type="checkbox"/> <input type="checkbox"/>	A. Rescuer 1 – push patient’s shoulders up and support patient’s back and head with body B. Rescuer 2 – Gently pull on patient’s arms
3. POSITIONING AND LIFTING	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Rescuer 1 – Support patient in sitting position B. Rescuer 2 – Slip hands under the patient’s knees C. On command, rescuers stand simultaneously, lifting patient with proper body mechanics

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

SHIRT DRAG

PROCEDURES		CRITICAL SKILL
1. POSITIONING	<input type="checkbox"/>	A. Rescuer - Kneel at the head of the patient and place one hand under each of the shoulders
2. MOVING PATIENT	<input type="checkbox"/> <input type="checkbox"/>	A. Rescuer - Grasp shirt at the shoulder area B. Drag patient in a straight line (keep spine as straight as possible)

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

ESTABLISHING AIRWAY-SUSPECTED CERVICAL SPINE (NECK) INJURY

PROCEDURES		CRITICAL SKILL
1. STABILIZE HEAD	<input type="checkbox"/> <input type="checkbox"/>	A. Rescuer – Position at top of the victim’s head B. Restrain victim’s head and neck to avoid voluntary or involuntary movement/rotation of the neck
2. ESTABLISH AIRWAY	<input type="checkbox"/>	A. Use modified jaw thrust maneuver without causing over-extension of victim’s neck
3. CHECK FOR BREATHING	<input type="checkbox"/> <input type="checkbox"/>	A. Look, listen and feel for breathing (3-5 seconds) B. State that the victim is/is not breathing
4. MAINTAIN OPEN AIRWAY	<input type="checkbox"/>	A. Do not compromise suspected neck injury

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

SHOCK

PROCEDURES		CRITICAL SKILL
1. CHECK FOR AND SYMPTOMS OF SHOCK	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Check for pale (or bluish) skin (in victim with dark skin examine inside of mouth and nailbeds for bluish coloration. *B. Check for cool, clammy skin *C. Check for weakness
2. TREATMENT	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Keep victim lying down B. Cover with blanket to prevent loss of body heat and place a blanket under the patient. (Do not try to place blanket under patient with possible spinal injuries) C. Elevate according to injury *D. Reassure and calm the patient

Option 1: Elevate the lower extremities or foot end of the back board. This procedure is performed in most cases. Place the patient flat, face up and elevate the legs or foot end of the back board 8 to 12 inches. Do not elevate any limbs with possible fractures or pelvic injuries until they have been properly splinted. Remember to consider the mechanism of injury for every patient.

Option 2: Lay the patient flat, face up. This is the supine position, used for patients with a spinal injury and patients who have serious injuries to the extremities that have not been supported. If the patient is placed in this position, you must constantly be prepared for vomiting.

Option 3: Slightly raise the head and shoulders. This position should be used only for responsive patients with no spinal injuries, life threatening chest or abdominal injuries and only for patients having difficulty breathing, but who have an open airway. A semi-seated position can also be used for patients with a history of heart problems. It is not recommended for moderate to severe cases of shock. Be certain to keep the patient's head from tilting forward.

Note: Injuries requiring the injured side to be tilted or placed down may be done after patient has been properly secured to a back board if a back board is required.

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

IMMOBILIZATION - LONG SPINE BOARD (Backboard)

PROCEDURES

CRITICAL SKILL

1. MOVE THE PATIENT ONTO THE LONG SPINE BOARD	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. One First Aid Provider at the head must maintain in-line immobilization of the head and spine B. First Aid Provider at the head directs the movement of the patient C. Other First Aid Provider control movement of the rest of body D. Other First Aid Provider position themselves on same side E. Upon command of First Aid Provider at the head, roll patient onto side toward First Aid Providers F. Quickly assess posterior body, if not already done G. Place long spine board next to the patient with top of board beyond top of head H. Place patient onto the board at command of the First Aid Provider at head while holding in-line immobilization using methods to limit spinal movement I. Slide patient into proper position using smooth coordinated moves keeping spine in alignment
2. PAD VOIDS BETWEEN PATIENT AND LONG SPINE BOARD	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Select and use appropriate padding B. Place padding as needed under the head C. Place padding as needed under torso
3. IMMOBILIZE BODY TO THE LONG SPINE BOARD	<input type="checkbox"/>	A. Strap and secure body to board ensuring spinal immobilization, beginning at shoulder and working toward feet
4. IMMOBILIZE HEAD TO THE LONG SPINE BOARD	<input type="checkbox"/> <input type="checkbox"/>	A. Using head set or place rolled towels on each side of head B. Tape and/or strap head securely to board, ensuring cervical spine immobilization
5. REASSESS	<input type="checkbox"/> <input type="checkbox"/>	*A. Reassess PMS (Pulse, Motor, Sensory) *B. Assess patient response and level of comfort

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

IMMOBILIZATION OF CERVICAL SPINE

PROCEDURES		CRITICAL SKILL
1. ESTABLISH AND IN-LINE IMMOBILIZATION	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Place head in a neutral, in-line position unless patient complains of pain or the head is not easily moved into position B. Place head in alignment with spine C. Maintain constant manual in-line immobilization until the patient is properly secured to a backboard with head immobilized
2. ASSESS PMS	<input type="checkbox"/>	*A. Assess PMS in all extremities: <ul style="list-style-type: none"> • Pulse • Motor function • Sensory function
3. ASSESS CERVICAL REGION AND NECK	<input type="checkbox"/>	*A. Inspect and palpate for injuries or signs of injuries using: DOTS acronym B. Remove clothing or jewelry as necessary
4. BANDAGE ANY WOUND	<input type="checkbox"/>	A. Any neck wounds
5. APPLY CERVICAL SPINE IMMOBILIZATION	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Apply properly sized collar or manual immobilization <u>One piece C-collar</u> <ul style="list-style-type: none"> A. Select proper sized collar B. Apply collar C. Ensure that patient's head is not twisted during application D. Ensure airway is open after placement <u>Two piece C-collar</u> <ul style="list-style-type: none"> A. Select proper sized collar B. Apply rear section to back of neck C. Center rigid support on spine D. Apply front section (overlaps rear section) E. Ensure chin rests in chin cavity F. Secure collar with Velcro straps G. Ensure airway is open after placement

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

6. SECURE HEAD TO APPROPRIATE IMMOBILIZATION DEVICE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Immobilize patient to appropriate immobilization device B. Use head set or place rolled blankets or towels on each side of head C. Tape head securely to appropriate immobilization Device
7. REASSESS	<input type="checkbox"/> <input type="checkbox"/>	*A. Reassess PMS *B. Assess patient response and level of comfort

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

BURNS

PROCEDURES

CRITICAL SKILLS

1. DETERMINE BURN TYPE	<input type="checkbox"/>	*A. Determine type <ul style="list-style-type: none"> ▪ Thermal ▪ Chemical ▪ Electrical
2. DETERMINE BODY SURFACE AREA	<input type="checkbox"/>	*A. Determine Body Surface Area (BSA) using rule of nines
3. BURN CARE (All Types)	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Remove patient from source of burn and prevent further contamination B. Consider the type of burn and stopping the burning process initially with water or saline if appropriate *C. Cut off smoldering clothing *D. Remove jewelry *E. Continually monitor the airway for evidence of closure F. Cover the burned area with a dry sterile dressing G. Do not use any type of ointment, lotion or antiseptic H. Do not break blisters I. Ensure patient does not get hypothermic
4. CARE FOR CHEMICAL BURNS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Protect yourself from exposure to hazardous materials B. Wear gloves, eye protection, and respiratory protection C. Brush off dry powders *D. Consider to flushing with large amounts of water *E. Continue flushing the contaminated area while en route to the receiving facility *F. Use caution not to contaminate uninjured areas when flushing

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

5. CARE FOR ELECTRICAL BURNS	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<p>*A. Ensure safety before removing patient from the electrical source</p> <p>*B. If the patient is still in contact with the electrical source or you are unsure, do not approach or touch the patient, contact power company</p> <p>*C. Monitor the patient closely for respiratory and cardiac arrest</p> <p>D. Treat the soft tissue injuries associated with the burn</p> <p>*E. Look for both an entrance and exit wound</p>
6. REASSESS	<input type="checkbox"/>	<p>*A. Reassess level of consciousness, respiratory status, and patient response</p>

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

EARLY OR SUPERFICIAL FROSTBITE

PROCEDURES		CRITICAL SKILLS
1. ASSESS FOR FROSTBITE AND COLD INJURIES	<input type="checkbox"/>	A. Patient exhibits signs and symptoms of frostbite or cold injuries
2. ASSESS FOR EARLY OR SUPERFICIAL FROSTBITE	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. Blanching of the skin – palpitation of the skin in which normal color does not return B. Loss of feeling and sensation in the injured area C. Skin remains soft D. If re-warmed, patient will feel a tingling sensation
3. TREAT EARLY OR SUPERFICIAL INJURY	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Remove the patient from the environment B. Protect the cold injured extremity from further injury *C. Remove wet or restrictive clothing D. Do not rub or massage E. Do not re-expose to the cold
4. REASSESS	<input type="checkbox"/>	*A. Reassess level of consciousness, respiratory status and patient response

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

LATE OR DEEP COLD INJURY

PROCEDURES		CRITICAL SKILLS
1. ASSESS FOR FROSTBITE AND COLD INJURIES	<input type="checkbox"/>	A. Patient exhibits signs and symptoms of frostbite or cold injuries
2. ASSESS FOR LATE OR DEEP COLD INJURY	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	A. White, waxy skin B. Firm to frozen feeling upon palpitation C. If thawed or partially thawed, the skin may appear flushed with areas of purple and blanching or mottled and cyanotic D. Swelling may be present E. Blisters may be present
3. TREAT LATE OR DEEP COLD INJURY	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Remove the patient from the environment B. Protect the cold injured extremity from further injury *C. Remove wet or restrictive clothing D. Remove jewelry E. Cover with dry clothing or dressings F. Do not: <ul style="list-style-type: none"> ▪ Break blisters ▪ Rub or massage area ▪ Apply heat ▪ Re-warm ▪ Allow the patient to walk on the affected extremity
4. REASSESS	<input type="checkbox"/>	*A. Reassess level of consciousness, respiratory status and patient response

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state "DOTS" when making their checks.

* Teams may use the acronym "PMS" when checking pulse, motor function and sensory function

MILD HYPERTHERMIA (HEAT)

PROCEDURES		CRITICAL SKILL
1. ASSESS FOR HYPERTHERMIA	<input type="checkbox"/>	A. Patient exhibits signs and symptoms of hyperthermia: <ul style="list-style-type: none"> ▪ Redness ▪ Muscular cramps ▪ Weakness or exhaustion ▪ Rapid heart rate ▪ Dizziness or faintness ▪ Altered mental status to unresponsive
2. PREVIOUS INTERVENTIONS	<input type="checkbox"/>	A. Inquire about previous interventions attempted
3. ASSESS FOR MILD HYPERTHERMIA (HEAT EXHAUSTION)	<input type="checkbox"/>	*A. Check skin for: <ul style="list-style-type: none"> ▪ Normal to cool temperature ▪ Pale ▪ Moist
4. TREATMENT FOR MILD HYPERTHERMIA	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Remove patient from hot environment and place in a cool environment *B. Loosen or remove clothing *C. Cool patient by fanning D. Put in supine position with legs elevated E. Offer drinking water if patient is responsive and not nauseated F. If the patient is unresponsive or is vomiting, transport to the hospital
5. REASSESS	<input type="checkbox"/>	*A. Reassess level of consciousness, respiratory status and patient response

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function

SEVERE HYPERTHERMIA

PROCEDURES

CRITICAL SKILL

1. ASSESS FOR HYPERTHERMIA	<input type="checkbox"/>	A. Patient exhibits signs and symptoms of hyperthermia: <ul style="list-style-type: none"> ▪ Redness ▪ Muscular cramps ▪ Weakness or exhaustion ▪ Rapid heart rate ▪ Dizziness or faintness ▪ Altered mental status to unresponsive
2. PREVIOUS INTERVENTIONS	<input type="checkbox"/>	A. Inquire about previous interventions attempted
3. ASSESS FOR SEVERE HYPERTHERMIA (HEAT STROKE)	<input type="checkbox"/>	*A. Check skin for: <ul style="list-style-type: none"> ▪ Hot temperature ▪ Red ▪ Dry or moist
4. TREATMENT FOR SEVERE HYPERTHERMIA	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	*A. Place patient in a cool environment *B. Remove clothing *C. Wet patient skin by applying water from sponge or wet towels and fan *D. Put in supine position with legs elevated *E. Offer drinking water if patient is responsive and not nauseated *F. Apply cool packs to neck, groin and armpits *G. Transport immediately
5. REASSESS	<input type="checkbox"/>	*A. Reassess level of consciousness, respiratory status and patient response

NOTE: Each critical skill identified with an asterisk (*) shall be clearly verbalized by the team as it is being conducted. After initially stating what DOTS stands for, the team may simply state “DOTS” when making their checks.

* Teams may use the acronym “PMS” when checking pulse, motor function and sensory function